

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51508537>

DMRT1 prevents female reprogramming in the postnatal mammalian testis

Article in *Nature* · August 2011

DOI: 10.1038/nature10239 · Source: PubMed

CITATIONS

302

READS

201

6 authors, including:



Clint Matson

Gilead Sciences

18 PUBLICATIONS 1,064 CITATIONS

[SEE PROFILE](#)



Mark W Murphy

University of Minnesota Twin Cities

26 PUBLICATIONS 1,370 CITATIONS

[SEE PROFILE](#)



Michael Griswold

Washington State University

346 PUBLICATIONS 15,294 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Human Testis and Spermatogenesis [View project](#)



Cell to cell interactions [View project](#)

DMRT1 prevents female reprogramming in the postnatal mammalian testis

Clinton K. Matson^{1,2}, Mark W. Murphy¹, Aaron L. Sarver³, Michael D. Griswold⁴, Vivian J. Bardwell^{1,2,3} & David Zarkower^{1,2,3}

Sex in mammals is determined in the fetal gonad by the presence or absence of the Y chromosome gene *Sry*, which controls whether bipotential precursor cells differentiate into testicular Sertoli cells or ovarian granulosa cells¹. This pivotal decision in a single gonadal cell type ultimately controls sexual differentiation throughout the body. Sex determination can be viewed as a battle for primacy in the fetal gonad between a male regulatory gene network in which *Sry* activates *Sox9* and a female network involving WNT/ β -catenin signalling². In females the primary sex-determining decision is not final: loss of the FOXL2 transcription factor in adult granulosa cells can reprogram granulosa cells into Sertoli cells². Here we show that sexual fate is also surprisingly labile in the testis: loss of the DMRT1 transcription factor³ in mouse Sertoli cells, even in adults, activates *Foxl2* and reprograms Sertoli cells into granulosa cells. In this environment, theca cells form, oestrogen is produced and germ cells appear feminized. Thus *Dmrt1* is essential to maintain mammalian testis determination, and competing regulatory networks maintain gonadal sex long after the fetal choice between male and female. *Dmrt1* and *Foxl2* are conserved throughout vertebrates^{4,5} and *Dmrt1*-related sexual regulators are conserved throughout metazoans³. Antagonism between *Dmrt1* and *Foxl2* for control of gonadal sex may therefore extend beyond mammals. Reprogramming due to loss of *Dmrt1* also may help explain the aetiology of human syndromes linked to *DMRT1*, including disorders of sexual differentiation⁶ and testicular cancer⁷.

Human chromosome 9p deletions removing *DMRT1* are associated with XY male-to-female sex reversal, and *Dmrt1* homologues determine sex in several non-mammalian vertebrates^{8–10}. In mice, *Dmrt1* is expressed and required in both germ cells and Sertoli cells of the testis^{11–13}. XY *Dmrt1*-null mutant mice are born as males with testes, although these gonads later undergo abnormal differentiation¹⁴; hence the role of *Dmrt1* in mammalian sex determination has been unclear (for overview of mammalian sex determination see Supplementary Fig. 1). Here we examine *Dmrt1* mutant testes during postnatal development, asking whether loss of *Dmrt1* causes postnatal feminization in mice.

We first examined gonads of *Dmrt1*-null mutant males (*Dmrt1*^{-/-}) for the presence of FOXL2, a female-specific transcription factor expressed in granulosa cells and theca cells^{15,16}, the two somatic cell types of the ovarian follicle (Fig. 1a). Four weeks after birth, abundant FOXL2-positive cells were present within mutant seminiferous tubules (Fig. 1b), which in control testes contain only germ cells and Sertoli cells (Fig. 1c). To establish the origin of the FOXL2-positive cells, we deleted *Dmrt1* either in germ cells (using *Nanos3-cre*) or in Sertoli cells (using *Dhh-cre* or *Sf1-cre*) (Supplementary Fig. 2a–l and Supplementary Table 1). Loss of *Dmrt1* in fetal Sertoli cells (*SCDmrt1KO*) but not in fetal germ cells (*GCDmrt1KO*) induced FOXL2 expression (Fig. 1d–f). *SCDmrt1KO* gonads retained small numbers of germ cells, which appeared to arrest in meiotic prophase on the basis of SYCP3 localization (Supplementary Fig. 3). These results demonstrate that DMRT1

expression in Sertoli cells prevents FOXL2 expression and suggest that *Dmrt1* mutant testes become feminized during the first postnatal month.

Next we examined the timing of FOXL2 induction. At postnatal day (P)7, *SCDmrt1KO* testes had seminiferous tubules in which all Sertoli cells expressed SOX9 normally (Supplementary Fig. 2m–r), but at P14




Figure 1 | DMRT1 maintains SOX9 and suppresses FOXL2 expression in postnatal Sertoli cells. **a–c**, FOXL2 expression detected by immunofluorescence in adult (Ad.) granulosa and theca cells of control ovary (**a**) and intratubular cells of *Dmrt1*-null testis at P28 (**b**), but not in control testis (**c**). DAPI, 4',6-diamidino-2-phenylindole. **d–f**, FOXL2 is robustly expressed when *Dmrt1* is mutated in fetal Sertoli cells with *Dhh-cre* (**d**) or *Sf1-cre* (**e**) but not when *Dmrt1* is mutated in fetal germ cells with *Nanos3-cre* (**f**). **g–o**, Timing of FOXL2 expression. FOXL2 is absent from control testis at P14 (**g–i**). Cells expressing FOXL2 or FOXL2 and SOX9 (arrowheads) are present in *SCDmrt1KO* testis at P14 (**j–l**). FOXL2-positive cells are abundant in *SCDmrt1KO* testis at P28 and most cells no longer express SOX9 (**m–o**). Scale bars, 20 μ m.

¹Developmental Biology Center and Department of Genetics, Cell Biology, and Development, University of Minnesota, Minneapolis 55455, Minnesota, USA. ²Molecular, Cellular, Developmental Biology, and Genetics Graduate Program, University of Minnesota, Minneapolis 55455, Minnesota, USA. ³University of Minnesota Masonic Cancer Center, Minneapolis, Minnesota 55455, USA. ⁴School of Molecular Biosciences, Washington State University, Pullman, Washington 99164, USA.





Figure 2 | Sertoli-to-granulosa transdifferentiation in the adult testis. **a–h**, Expression of FOXL2 and SOX9 one month after tamoxifen (TX) injection into *Dmrt1^{fllox/fllox}* adult males (8 weeks and older) carrying inducible ubiquitous *cre* transgene *UBC-cre/ERT2*. **a, b**, Sertoli cells in control testis express SOX9 but not FOXL2. **c–f**, Mutant testis has Sertoli-like cells expressing SOX9 or SOX9 and FOXL2 (**d**, inset) and granulosa-like cells expressing only FOXL2

some intratubular cells co-expressed SOX9 and FOXL2 or lacked SOX9 and strongly expressed FOXL2 (Fig. 1g–l). By P28 few SOX9-positive cells remained and most intratubular cells strongly expressed FOXL2 (Fig. 1m–o). Histological analysis of mutant gonads is shown in Supplementary Fig. 4. These results show that fetal loss of *Dmrt1* causes postnatal Sertoli cells to lose the male-promoting SOX9 and instead express the female-promoting FOXL2.

Loss of *Foxl2* in the adult ovary can lead to transdifferentiation of granulosa cells to Sertoli cells², so we asked whether loss of *Dmrt1* in the adult testis activates *Foxl2* and causes the reciprocal sex transformation, from Sertoli to granulosa. Indeed, one month after deletion of *Dmrt1* in adult males (using a tamoxifen-inducible *cre* transgene), we observed cells with typical Sertoli cell features including tripartite nucleoli but expressing both SOX9 and FOXL2 (Fig. 2a–d), as well as



(f, inset). **g, h**, FOXL2-positive cells in control ovary have DAPI morphology similar to FOXL2 single-positive cells of mutant testis. FOXL2-positive cells in mutant testis resemble granulosa cells: they lack the tripartite nucleoli of Sertoli cells, have smaller and more rounded nuclei, and have more punctate DAPI staining. *UBC-cre/ERT2* also deletes *Dmrt1* in germ cells, causing precocious meiosis¹²; after one month germ cells are nearly absent. Scale bars, 20 μm.

cells with typical granulosa cell nuclear morphology that lacked SOX9 and strongly expressed FOXL2 (Fig. 2e–h). Thus antagonism between DMRT1 and FOXL2 continues into adulthood and Sertoli cell fate remains plastic even after terminal differentiation.

To evaluate further the transformation of mutant gonads, we compared the messenger RNA profile of control and mutant P28 testes; 5,030 mRNAs were expressed >8-fold differently across this data set or a data set comparing testis to 21 other tissues including ovary (Supplementary Fig. 5a). We calculated Pearson correlation coefficients for expression of these 5,030 mRNAs in mutant gonads relative to each tissue and found that the mutant gonad most closely resembled ovary (Supplementary Fig. 5b; average $R = 0.75$). Many mRNAs with decreased expression in mutant gonads also were low in other tissues, probably reflecting a lack of male germ cells, which comprise much of the testis mass. Also, some mRNAs that were increased in mutant gonads were increased in other tissues. Therefore, to specifically evaluate ovary-enriched mRNAs, we used bioGPS (<http://www.biogps.gnf.org>; see Supplementary Information) to identify 65 mRNAs with expression closely correlated to *Foxl2* and then compared their expression in ovary relative to the other 21 tissues (Fig. 3a and Supplementary Fig. 6). This comparison confirmed that these mRNAs are highly ovary enriched. About 40% were increased in mutant gonads relative to control testes; about 80% of the remainder were oocyte enriched. Thus loss of *Dmrt1* causes large changes in mRNA expression, including induction of multiple ovary-enriched mRNAs. mRNA profiling of *Dmrt1* mutant gonads perinatally and at P9 did not reveal

Figure 3 | Feminization of *SCDmrt1KO* XY gonads. **a**, Expression of ovary-enriched mRNAs with expression profiles similar to *Foxl2* (see Supplementary Information). mRNAs labelled 'somatic' were enriched in ovarian somatic cells; those labelled 'oocyte' were enriched in female germ cells. See Supplementary Fig. 6 for higher resolution image. **B**, marrow, bone marrow; Saliv. gland, salivary gland; Sem. vesicle, seminal vesicle; Sm. intestine, small intestine. **b–d**, Immunohistochemistry detection of CYP19A1/aromatase expression in follicles of control adult ovary (**a**) and in adult XY *SCDmrt1KO* gonad (**b**), but only in interstitial Leydig cells of control testis (**c**). Arrows indicate aromatase-positive granulosa cells in ovary and mutant gonad and negative Sertoli cell in control testis. Scale bars, 50 μm. **e–g**, Immunofluorescence detection of SMA and FOXL2. Ovarian theca cells (**e**, inset) are elongated cells expressing both proteins; similar cells are present in mutant gonads (**f**); peritubular myoid cells in control testes express SMA and not FOXL2 (**g**). Scale bars, 20 μm. **h–j**, Immunofluorescence detection of cells coexpressing FOXL2 in the nucleus and steroidogenic enzyme CYP11A1/SCC at high levels in the cytoplasm in control ovary (**h**) and XY *SCDmrt1KO* gonads (**i**). SCC-positive cells in control testis (**j**) are interstitial Leydig cells. Mutant gonads were *SCDmrt1KO(Dhh)*. All tissues were from adult mice. Scale bars, 20 μm.

apparent feminization^{17,18}, consistent with the observation that FOXL2 expression starts at ~P14.

Further analysis of the mRNA profiling data identified highly increased expression (>5-fold, $P < 0.001$) of many mRNAs expressed in granulosa cells and required for ovarian development or function. These included *Foxl2*, *Nr5a2* (also known as *Lrh1*), *Wnt4*, LH receptor (*Lhcgr*), prolactin receptor (*Prlr*), FSH receptor (*Fshr*), follistatin (*Fst*), *Sfrp4*, *Igfbp5*, *Inhbb*, *Inha* and *Lnfg* (Supplementary Table 2). *Foxl2os*, a noncoding RNA transcribed from the opposite strand of the *Foxl2* coding region, also was highly overexpressed and has been suggested as a positive regulator of *Foxl2* (ref. 19). We confirmed increased expression in mutant gonads of LRH1, a transcription factor expressed only in granulosa cells within the ovary²⁰ and absent from the testis (Supplementary Fig. 7a–f). *Nr5a2* is probably a direct target of DMRT1 regulation, based on binding of DMRT1 to its promoter proximal sequences *in vivo* (Supplementary Fig. 7g). On the basis of mRNA and protein expression data and changes in cellular morphology, we conclude that loss of *Dmrt1* in testes reprograms Sertoli cells into granulosa cells.

Granulosa cells produce oestrogens, which are essential for ovarian development in many vertebrates; in mammals, oestrogen signalling also acts with FOXL2 to repress *Sox9* transcription in adult granulosa cells². HSD17 β 1 and CYP19A1/aromatase are enzymes critical for oestrogen synthesis, and mRNAs for both enzymes were increased in mutant gonads (Supplementary Fig. 8). Aromatase protein is robustly expressed in granulosa cells and was strongly expressed in mutant gonads (Fig. 3b–d). Consistent with these enzyme changes, oestradiol was raised in the serum of adult mutants relative to control adult males (Supplementary Information). Although expression of the androgenic enzyme *Hsd17 β 3* was not affected in mutant gonads (Supplementary Fig. 8), androgen levels were reduced based on severely decreased seminal vesicle weight, a sensitive indicator of androgen activity (350 ± 52 mg versus 182 ± 36 mg; $n = 3$, $P = 0.01$).

Theca cells are induced during follicle growth in the ovary, probably in response to granulosa cell signals²¹, and together with granulosa cells and oocytes they comprise the functional unit of the ovary. Because mutant gonads contained apparently functional granulosa cells, we asked whether theca cells also formed. Theca cells have spindle-shaped nuclei and express both FOXL2 and smooth muscle actin (SMA) (Fig. 3e). Adult mutant gonads contained cells closely resembling theca cells and expressing both proteins (Fig. 3f). The theca-like cells probably derive either from granulosa cells or peritubular myoid cells (which also are elongated and express SMA; Fig. 3g). However, as seminiferous tubule integrity was lost before formation of these cells (Fig. 3f and Supplementary Fig. 9), they could potentially derive from interstitial cells that invaded the tubule remnants. We also observed intratubular cells strongly expressing the steroidogenic enzyme SCC (Fig. 3h–j); these cells resembled luteinized granulosa cells of the ovary (Fig. 3h), suggesting that granulosa cells in the mutant gonad are responsive to gonadotropins. We therefore tested the effect of exogenous gonadotropin stimulation; treated mutants, but not controls, had additional luteinized granulosa cells and germ cells with oocyte-like nuclear morphology that expressed the oocyte-specific proteins MATER and ZP2 (Supplementary Fig. 10). This result indicates that both somatic cells and germ cells are feminized in mutant gonads.

The preceding results indicate that DMRT1 is essential for postnatal sex maintenance. DMRT1 is a sequence-specific transcriptional regulator, capable of activating or repressing transcription of target genes^{18,22}. To help find targets of DMRT1 regulation with potential roles in sex maintenance we examined expression of known fetal sex-determining genes in mutant gonads at P28 by quantitative polymerase chain reaction with reverse transcription (qRT–PCR; Fig. 4a). Among masculinizing genes, *Ptgdr*, *Sox9* and *Sox8*, which acts redundantly with *Sox9* (refs 23, 24), were reduced. Among feminizing genes, *Foxl2*, *Esr1*, *Esr2*, *Wnt4* and *Rspo1* were increased. We assayed binding of DMRT1 to DNA of P28 testes by quantitative chromatin immunoprecipitation




Figure 4 | DMRT1 regulation of postnatal gene expression. **a**, qRT–PCR analysis of sex-determining genes at P28. Significance of expression changes is indicated (Student's *t*-test). Mutant gonads were *SCDmrt1KO*(*Sfl*); *SCDmrt1KO*(*Dhh*) mutant gonads and equivalent expression changes. **b**, qChIP analysis of DMRT1 DNA binding in P28 testes. Significance of enrichment relative to *B2m* (Student's *t*-test) is shown. **c**, Model for regulation by postnatal sex maintenance by DMRT1. Proposed direct regulation based on ChIP and mRNA expression data are indicated by solid lines; indirect or potential regulation is indicated by dashed lines. Model adapted from ref. 2.

(qChIP), guided by genome-wide ChIP data from P9 testes (ChIP-chip¹⁸ and ChIP-seq (unpublished data)). DMRT1 bound both upstream and downstream of *Sox9* and upstream of *Sox8*, and bound weakly near *Ptgdr*. DMRT1 bound strongly near *Foxl2*, *Esr1*, *Esr2*, *Wnt4* and *Rspo1* (Fig. 4b). All of the DMRT1-associated regions contained at least one close match to the DMRT1 DNA-binding consensus^{18,22}.

On the basis of mRNA and protein expression data and ChIP analysis, we propose a model for postnatal sex maintenance (Fig. 4b) in which DMRT1 maintains male fates by repressing multiple female-promoting genes and activating male-promoting genes. *Sox9* is dispensable for testis differentiation after sex determination^{24,25}, suggesting that other critical male regulators remain to be found; *Sox8* is a clear candidate based on its redundancy with *Sox9* (refs 23, 24). We find that DMRT1 represses *Foxl2*, which is known to maintain postnatal ovarian fate. FOXL2 also represses *Dmrt1* (ref. 2); thus antagonism between these sex-specific transcriptional regulators may be central to sex maintenance in both sexes throughout reproductive life. *Wnt4* and *Rspo1* also are prime candidates for postnatal sex maintenance based on their requirement in ovarian determination in the fetus^{26,27}. Indeed, P28 mutant gonads had increased nuclear β -catenin in somatic cells, as in ovaries, but control testes did not, indicating active WNT/ β -catenin signalling in the mutant gonads (Supplementary Fig. 11). Functional analysis of *Wnt4*, *Rspo1* and other known fetal sex regulators will be important to establish their roles in sex maintenance.

The analysis presented here demonstrates that deletion of *Dmrt1* during fetal development induces postnatal feminization of the testis, causing male-to-female primary sex reversal. Moreover, deletion of

Dmrt1 in adults can reprogram differentiated Sertoli cells into apparent granulosa cells. Why *Dmrt1* mutants are feminized only after birth remains unclear. Another male-promoting gene may act redundantly with *Dmrt1* before P14, masking its function; alternatively, the testis may lack potential feminizing activity from genes such as *Foxl2* before P14. Another puzzle is that *Dmrt1* mutant mice are born male, whereas human 9p deletions removing *DMRT1* can cause XY feminization at birth. The human sex reversal may reflect failure to maintain male sex determination, and the longer human gestation may permit testis-to-ovary reprogramming before birth. Alternatively, human testes may have potential feminizing activity earlier or may lack masculinizing genes redundant with *DMRT1*. Our results may provide insights into the aetiology of human gonadal disorders, including gonadoblastoma and granulosa cell tumours of the testis. Moreover, because many genes implicated in this study are evolutionarily conserved, similar mechanisms may control adult sex switching in fish and may maintain sexual fate in the adult gonads of other vertebrates or even in other phyla.

METHODS SUMMARY

Mouse breeding. *Dmrt1* mutant and control males were generated as described¹²; tissue-specific Cre recombinase strains are in Supplementary Table 1. Adult wild-type or *Dmrt1*^{flax/flax} females were used as controls. Mice were mixed C57BL/6J, 129S1 and FVB genetic background. Protocols were approved by the Institutional Animal Care and Use Committee.

Immunofluorescence and immunohistochemistry. Immunofluorescence and immunohistochemistry were performed as described¹². Antibodies are listed in Supplementary Table 3. Analyses included at least two biological replicates.

Tamoxifen treatment. Tamoxifen-inducible deletion of *Dmrt1* in adult males was as described¹². Testes were harvested one to two months after treatment.

mRNA expression analysis. mRNA expression profiling and data analysis were as described¹³ except total testis RNA was isolated from 4-week-old mice using TRIzol reagent (Invitrogen no. 15596-026). Additional detail is in Supplementary Methods.

qRT-PCR. qRT-PCR was as described¹². qRT-PCR primers are listed in Supplementary Table 4.

ChIP. ChIP followed by either microarray (ChIP-chip) or qPCR analysis (qChIP) were as described¹⁸. Gene-specific primers used for qChIP are in Supplementary Table 4.

Full Methods and any associated references are available in the online version of the paper at www.nature.com/nature.

Received 11 February; accepted 27 May 2011.

Published online 20 July 2011.

- Koopman, P., Gubbay, J., Vivian, N., Goodfellow, P. & Lovell-Badge, R. Male development of chromosomally female mice transgenic for *Sy*. *Nature* **351**, 117–121 (1991).
- Uhlenhaut, N. H. *et al.* Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell* **139**, 1130–1142 (2009).
- Raymond, C. S. *et al.* Evidence for evolutionary conservation of sex-determining genes. *Nature* **391**, 691–695 (1998).
- Loffler, K. A., Zarkower, D. & Koopman, P. Etiology of ovarian failure in blepharophimosis ptosis epicanthus inversus syndrome: *FOXL2* is a conserved, early-acting gene in vertebrate ovarian development. *Endocrinology* **144**, 3237–3243 (2003).
- Raymond, C. S., Kettlewell, J. R., Hirsch, B., Bardwell, V. J. & Zarkower, D. Expression of *Dmrt1* in the genital ridge of mouse and chicken embryos suggests a role in vertebrate sexual development. *Dev. Biol.* **215**, 208–220 (1999).
- Tannour-Louet, M. *et al.* Identification of *de novo* copy number variants associated with human disorders of sexual development. *PLoS ONE* **5**, e15392 (2010).
- Turnbull, C. *et al.* Variants near *DMRT1*, *TERT* and *ATF7IP* are associated with testicular germ cell cancer. *Nature Genet.* **42**, 604–607 (2010).

- Yoshimoto, S. *et al.* A W-linked DM-domain gene, DM-W, participates in primary ovary development in *Xenopus laevis*. *Proc. Natl Acad. Sci. USA* **105**, 2469–2474 (2008).
- Smith, C. A. *et al.* The avian Z-linked gene *DMRT1* is required for male sex determination in the chicken. *Nature* **461**, 267–271 (2009).
- Matsuda, M. *et al.* *DMY* is a Y-specific DM-domain gene required for male development in the medaka fish. *Nature* **417**, 559–563 (2002).
- Kim, S., Bardwell, V. J. & Zarkower, D. Cell type-autonomous and non-autonomous requirements for *Dmrt1* in postnatal testis differentiation. *Dev. Biol.* **307**, 314–327 (2007).
- Matson, C. K. *et al.* The mammalian doublesex homolog *DMRT1* is a transcriptional gatekeeper that controls the mitosis versus meiosis decision in male germ cells. *Dev. Cell* **19**, 612–624 (2010).
- Krentz, A. D. *et al.* The DM domain protein *DMRT1* is a dose-sensitive regulator of fetal germ cell proliferation and pluripotency. *Proc. Natl Acad. Sci. USA* **106**, 22323–22328 (2009).
- Raymond, C. S., Murphy, M. W., O'Sullivan, M. G., Bardwell, V. J. & Zarkower, D. *Dmrt1*, a gene related to worm and fly sexual regulators, is required for mammalian testis differentiation. *Genes Dev.* **14**, 2587–2595 (2000).
- Schmidt, D. *et al.* The murine winged-helix transcription factor *Foxl2* is required for granulosa cell differentiation and ovary maintenance. *Development* **131**, 933–942 (2004).
- Uda, M. *et al.* *Foxl2* disruption causes mouse ovarian failure by pervasive blockage of follicle development. *Hum. Mol. Genet.* **13**, 1171–1181 (2004).
- Fahrioglu, U., Murphy, M. W., Zarkower, D. & Bardwell, V. J. mRNA expression analysis and the molecular basis of neonatal testis defects in *Dmrt1* mutant mice. *Sex. Dev.* **1**, 42–58 (2007).
- Murphy, M. W. *et al.* Genome-wide analysis of DNA binding and transcriptional regulation by the mammalian Doublesex homolog *DMRT1* in the juvenile testis. *Proc. Natl Acad. Sci. USA* **107**, 13360–13365 (2010).
- Cocquet, J., Pannetier, M., Fellous, M. & Veitia, R. A. Sense and antisense *Foxl2* transcripts in mouse. *Genomics* **85**, 531–541 (2005).
- Duggavathi, R. *et al.* Liver receptor homolog 1 is essential for ovulation. *Genes Dev.* **22**, 1871–1876 (2008).
- Orisaka, M., Tajima, K., Tsang, B. K. & Kotsuji, F. Oocyte-granulosa-theca cell interactions during preantral follicular development. *J. Ovarian Res.* **2**, 9 (2009).
- Murphy, M. W., Zarkower, D. & Bardwell, V. J. Vertebrate DM domain proteins bind similar DNA sequences and can heterodimerize on DNA. *BMC Mol. Biol.* **8**, 58 (2007).
- Chaboissier, M. C. *et al.* Functional analysis of *Sox8* and *Sox9* during sex determination in the mouse. *Development* **131**, 1891–1901 (2004).
- Barrionuevo, F. *et al.* Testis cord differentiation after the sex determination stage is independent of *Sox9* but fails in the combined absence of *Sox9* and *Sox8*. *Dev. Biol.* **327**, 301–312 (2009).
- Chang, H. *et al.* *Wt1* negatively regulates β -catenin signaling during testis development. *Development* **135**, 1875–1885 (2008).
- Vainio, S., Heikkilä, M., Kispert, A., Chin, N. & McMahon, A. P. Female development in mammals is regulated by *Wnt-4* signalling. *Nature* **397**, 405–409 (1999).
- Parma, P. *et al.* *R-spondin1* is essential in sex determination, skin differentiation and malignancy. *Nature Genet.* **38**, 1304–1309 (2006).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Acknowledgements We thank M. Treier for helpful discussion, K. Hatzl, A. Minkina, A. Peterson, the University of Minnesota Mouse Genetics Laboratory and C. Small for technical assistance, J. Dean, R. Veitia and K.-i. Morohashi for antibodies, D. Greenstein and A. M. Weber-Main for comments on the manuscript, C. Manivel for histology expertise, M. Steffes and D. Gabrielson for oestradiol analysis, and the University of Minnesota Supercomputing Institute for computational resources. This work was funded by the NIH (GM59152), the Minnesota Medical Foundation, and a predoctoral fellowship from the NSF (to C.K.M.).

Author Contributions C.K.M. performed mouse breeding and analysis of protein and mRNA expression; M.W.M. performed ChIP analysis; A.L.S. performed bioinformatic analysis; C.K.M., D.Z. and V.J.B. designed the study, analysed data, and wrote the paper; M.D.G. provided mRNA profiling expertise; all authors discussed the results and edited the paper.

Author Information mRNA expression profiling data have been deposited at the Gene Expression Omnibus under accession number GSE27261 and can be reviewed at <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?token=z1ctbiugsamymtc&acc=GSE27261>. Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at www.nature.com/nature. Correspondence and requests for materials should be addressed to D.Z. (zarko001@umn.edu) or V.J.B. (bardw001@umn.edu).

METHODS

Mouse breeding. Conditional *Dmrt1* mutant and control males were generated as described¹²; tissue-specific Cre recombinase strains are in Supplementary Table 1. Adult wild-type or *Dmrt1*^{fllox/fllox} females were used as controls. Mice were of mixed C57BL/6J, 129S1 and FVB genetic background. Protocols were approved by the University of Minnesota Institutional Animal Care and Use Committee.

Immunofluorescence and immunohistochemistry. Both immunofluorescence and immunohistochemistry were performed as described¹². Antibodies are listed in Supplementary Table 3. Analyses included a minimum of two biological replicates.

Tamoxifen treatment. Tamoxifen-inducible deletion of *Dmrt1* in adult males was performed as previously described¹². Testes were harvested one to two months after treatment.

mRNA expression analysis. mRNA expression profiling and data analysis were performed as described¹³ except total testis RNA was isolated from 4-week-old mice using TRIzol reagent (Invitrogen no. 15596-026). Affymetrix Mouse Genome 439 2.0 arrays were normalized by GC-RMA normalization²⁸ using GeneData Refiner. The Raw .cel files and the normalized data are deposited in the Gene Expression Omnibus (GEO)²⁹ under accession number GSE27261. GSE9954 was obtained from the GEO database. The arrays with the highest sample identification numbers were removed from the tissue data set to select 22 tissue types, each with three experimental replicates. When multiple probe sets were mapped to the same gene symbol, these values were averaged to obtain one value for each gene symbol. Direct Pearson correlation *R* values were calculated using all array data following reduction to gene symbols, and these values are shown in Fig. 2b.

Each experiment in our data set was divided by the average expression value from control testis tissue. GSE9954 data were separately divided by the average signal obtained from the GSE9954 testis samples. This was done separately for each data set to determine how samples from each data set differed from a baseline 'testis' expression state. Cluster 3.0 software³⁰ was used to: (1) log base 2 transform

the data; (2) filter the data set for genes that showed at least three observations with $\text{abs}(\text{val}) > 3$ (eightfold), which resulted in 5,030 genes passing the filter using both data sets combined; and (3) cluster the data on the gene axis using average linkage hierarchical clustering. The experimental axis was defined by order of decreasing correlation to the mutant testes calculated as described earlier. Javatreeview Software³¹ was used to generate heatmap images.

qRT-PCR. qRT-PCR was performed as described¹². qRT-PCR primers are listed in Supplementary Table 4.

Chromatin immunoprecipitation. ChIP followed by either microarray (ChIP-chip) or qPCR analysis (qChIP) were performed as described¹⁸. Gene-specific primers used for qChIP are in Supplementary Table 4.

Oestradiol assays. Serum oestradiol was assayed using a clinical electrochemiluminescence immunoassay (Roche Estradiol II, 03000079 122) according to manufacturer's instructions. Three of three males assayed had levels below the detection limit, whereas two of three females had measurable oestradiol (5.0 and 19.7 pg dl⁻¹). Two of three *SCDmrt1KO(Dhh)* mutant males had measurable oestradiol (5.6 and 21.2 pg dl⁻¹).

Gonadotropin treatment. Six-to-eight-week-old mutant males, control males and control females were treated with 5 units of pregnant mare serum by intraperitoneal injection and gonads were harvested 48 h later.

28. Wu, J., Irazarray, R. A., Gentleman, R., Martinez-Murillo, F. & Spencer, F. A model-based background adjustment for oligonucleotide expression arrays. *J. Am. Stat. Assoc.* **99**, 909–917 (2004).
29. Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res.* **30**, 207–210 (2002).
30. de Hoon, M. J., Imoto, S., Nolan, J. & Miyano, S. Open source clustering software. *Bioinformatics* **20**, 1453–1454 (2004).
31. Saldanha, A. J. Java Treeview—extensible visualization of microarray data. *Bioinformatics* **20**, 3246–3248 (2004).